

OptiPrep™ Application Sheet S59

Isolation of organelles from algae and fungi

- ◆ OptiPrep™ is a 60% (w/v) solution of iodixanol in water, density = 1.32 g/ml
- ◆ To access other Application Sheets referred to in the text: return to the 2020SMemapp file and select the appropriate S-number.
- ◆ Yeast organelles are covered in other Application Sheets; see Subcellular Membrane Index
- ◆ This Application Sheet summarizes all of the known OptiPrep™ applications for algae and fungi; a detailed methodology is only provided in a few instances.
- ◆ Some of the more recent papers are summarized in Section 7.

1. *Chlamydomonas reinhardtii*

Gradients that are used primarily for the isolation of acidocalcisomes also resolve mitochondria; see Application Sheet S50.

Membrane vesicles can be shed from flagella by shaking in 25 mM KCl, 5 mM MgSO₄, 1 mM DTT, 0.5 mM EDTA, 10 mM Hepes, pH 7.2 containing 0.1% NP40 (plus protease inhibitors). Axonemes are then removed by repeated centrifugation at 16,000 g for 10 min and the vesicles harvested from the supernatant at 228,000 g for 30 min. The crude vesicles in 1 ml of the medium are adjusted to 15% (w/v) iodixanol; underlaid with an equal column of 30% iodixanol and centrifuged at 431,000 g for approx 1 h. The vesicles band in the middle of the gradient [1].

In a study of the development of cilia in Wood and Rosenbaum in *Chlamydomonas reinhardtii* [2] sCytoplasmic extract was adjusted to 30% w/v iodixanol and overlaid by layers of 25%, 20%, 15%, 12%, 10%, and 5% and centrifuged for 3 hr at 200,000 g. The gradients were used to study the attachment of proteins to cytoplasmic membranes during flagellar development.

2. *Cladopeltis resinae*

The LMF in a routine buffered 0.25 M sucrose solution containing 1 mM EDTA was adjusted to 35% (w/v) iodixanol and a gradient formed by self-generation in near vertical NVT65 rotor centrifuged at 202,000 g for 4 h [3]. The organelle distribution is shown diagrammatically in Figure 1A; if the overlapping fractions of mitochondria and lysosomes are collected and adjusted to 25% iodixanol and re-centrifuged then a complete separation of these two organelles is obtained (Figure 1B).




Figure 1 Sequential separation of *Cladopeltis* organelles in two self-generated gradients: A, 35% iodixanol, B 25% iodixanol. For more information see text.

3. *Cyanidioschyzon merolae*

3.1 Microtubule-linked organelles

Cells were lysed in a microtubule stabilizing medium: 1.2 M sorbitol, 2 mM MgSO₄, 2 mM KCl, 1 mM DTT, 4 mM EGTA, 10 mM Tris-HCl, pH 7.2, containing, 50µM paclitaxel and 0.1% BSA, plus protease inhibitors [4,5]. The homogenate was mixed with an equal volume of 10% (w/v) iodixanol in 0.6 M sorbitol, 2 mM MgSO₄, 2 mM KCl, 1 mM DTT, 4 mM EGTA, 10 mM Tris-HCl, pH 7.2, containing, 10µM paclitaxel (plus protease inhibitors). A discontinuous gradient of 15%, 25% and 40% (w/v) iodixanol (8 ml, 8 ml and 4 ml respectively), in the same buffer, were overlaid with 15 ml of 300 g supernatant and centrifuged at 141,000 g for 30 min. The microtubule-linked organelles banded at the lowest interface.

3.2 Polyphosphate vacuoles

Yagisawa et al [6] isolated these polyphosphate-rich vacuoles from *Cyanidioschyzon merolae* and identified a number of important proteins. The discontinuous gradient used by the authors was an

unusual one, comprising three layer one of 27%, 62% and 80% (w/v) iodixanol. To achieve the latter two solutions, the authors first produced a very high density stock by evaporating the water from OptiPrep™. **For safety reasons this is not a procedure that the manufacturers of OptiPrep™ can recommend.** Since however the vacuoles banded at the 27%/62% iodixanol interface we suggest that the densest layer is omitted and the gradient formed from OptiPrep™ and 27% (w/v) iodixanol. To maintain the other solution properties used by the authors first prepare a diluent of 62% (w/w) sucrose solution containing 40 mM MgCl₂, 100 mM KCl, 400 mM Tris-HCl, pH 7.6. For the dense gradient solution add 1 ml of the diluent to 20 ml of OptiPrep™. From this prepare a 27% (w/v) iodixanol solution by diluting 2.7 vol. with 3.3 ml of 0.25 M sucrose, 2 mM MgCl₂, 5 mM KCl, 20 mM Tris-HCl, pH 7.6. Use the latter to resuspend the total membrane pellet. Layer 0.6 ml of this on to 2 ml of the dense solution and overlay with 1.4 ml of the lysis solution and centrifuge at 50,000 g for 1 h at 8°C. The vacuoles band at the lower interface.

- ◆ More recently, iodixanol gradients have been used to study peroxisomes from this algae [7].

4. *Cryptococcus neoformans* extracellular vesicles (EVs)

The glucuronoxylomannan of this fungus appears to be transported into the extracellular space in a membrane vesicle. This exocytic process is akin to that which occurs in mammalian cells and the shallow 6-18% (w/v) iodixanol gradients used to purify the EVs released from such cells have also been used for similar studies with *Cryptococcus neoformans*. The gradients were centrifuged at 250,000 g for 75 min [8]. A complex banding pattern of the glucuronoxylomannan and sterol suggest a heterogeneity of the budded vesicles. Discontinuous iodixanol gradients have been used by Wolf et al [9] comprising 3 ml of 30% and 2 ml each of 25%, 20%, 15% and 10% (w/v) iodixanol. The crude membrane vesicles in 5% (w/v) iodixanol were layered on top and centrifuged at 140 000 g, at 4°C, for 15 h. The peak of EVs occurred just below the mid-point of the gradient and interestingly the lightest ones were deficient in glucuronoxylomannan.

5. *Neurospora crassa*

Glyoxysomes have been prepared from an *N. crassa* post-nuclear supernatant (PNS) using a gradient similar [10] to that developed for mammalian peroxisomes [see Application Sheet S12](#). The following method is adapted from ref 10.

5.1 Solutions required

- A. OptiPrep™
- B. OptiPrep™ diluent: 0.25 M sucrose, 10 mM EDTA, 100 mM MOPS-NaOH pH 7.2, 1% (v/v) ethanol (see Box)
- C. 54% (w/v) Iodixanol stock solution: 9 vol. of Solution A + 1 vol. Solution B.
- D. Homogenization medium: 0.25 M sucrose, 1 mM EDTA, 10 mM MOPS-NaOH, pH 7.2, containing 0.1% (v/v) ethanol.
- E. Gradient solutions of 47% and 35% (w/v) iodixanol: dilute Solution C with Solution D.

Keep the following stock solutions at 4°C:	
1 M MOPS	20.9 g per 100 ml water
100 mM EDTA (Na ₂ •2H ₂ O)	3.72 g per 100 ml water
Solution B: Dissolve 17 g sucrose in 100 ml water; add 2 ml, 20 ml and 20 ml respectively of ethanol, EDTA stock and Mops stock; adjust to pH 7.2 with 1 M NaOH and make up to 200 ml.	
Solution D: Dissolve 17 g sucrose in 100 ml water; add 0.2 ml, 2 ml and 2 ml respectively of ethanol, EDTA stock and Mops stock; adjust to pH 7.2 with 1 M NaOH and make up to 200 ml.	

5.2 Ultracentrifuge rotor

Sorvall TH641 or Beckman SW41 or equivalent rotor with approx. 30 ml tubes

5.3 Protocol

1. Prepare an homogenate of *N. crassa* in Solution D.
2. Centrifuge the homogenate at 500 g for 5 min and repeat this three times to remove all of the abrasive powder and nuclei.
3. Prepare a crude organelle pellet was by centrifugation of the PNS at 25,000 g for 30 min and resuspend the pellet in Solution D.

4. Mix the resuspended pellet with an equal volume of the 47% (w/v) iodixanol solution.
5. Transfer 9-10 ml of the crude organelle suspension in 23.5% iodixanol and underlayer with 1-2 ml of the 35% iodixanol solution. If necessary top up the tubes to the required filling volume by layering Solution D on top.
6. Centrifuge at 110,000 g for 2 h. The glyoxysomes band at the interface between the two iodixanol solutions.

6. *Paracoccidioides brasiliensis*

The mitochondria and peroxisomes from the yeast phase of *Paracoccidioides brasiliensis* have been isolated using a gradient identical to that used for preparing mitochondria from *S. cerevisiae*.

Section 4 of Application Sheet S17 describes the discontinuous gradient that is used [11]; the densest band contains the peroxisomes. See also **Application Sheet S57** for more information of the isolation of peroxisomes from *S. cerevisiae*.

7. Recent publications

A CsCl gradient was used primarily to separate flagella, transition zones, and empty flagellar collars from *Chlamydomonas elegans*. To improve the purity of the transition zone fraction, it was diluted with buffer and harvested by centrifugation at approx 190,000 g_{av} for 20 min. The pellet was resuspended in buffer; adjusted to 30% iodixanol and made part of a discontinuous 60%, 30%, 20% (w/v) iodixanol step gradient. After centrifugation for 3 hr at approx 165,000 g_{av}. The transitional zones banded at the 60%/30% iodixanol interface, diluted. For more details see ref 12. Secretory vesicles from *Candida albicans* [13] were fractionated according to a method first described by Chang et al (see **Application Sheet S52**).

- ◆ Note the other **Application Sheets** devoted to algae and fungi:
- ◆ Algae: Acidocalcisomes (**S50**)
- ◆ Fungi: Extracellular vesicles (**S62**)

8. References

1. Huang, K., Diener, D.R., Mitchell, A., Pazour, G.J., Witman, G.B. and Rosenbaum, J.L. (2007) *Function and dynamics of PKD2 in Chlamydomonas reinhardtii flagella* J. Cell Biol., **179**, 501-514
2. Wood, C.R. and Rosenbaum, J.L. (2014) *Proteins of the ciliary axoneme are found on cytoplasmic membrane vesicles during growth of cilia* Curr. Biol., **24**, 1114-1120
3. Goswami, P. and Cooney, J.J. (1999) *Subcellular location of enzyme involved in oxidation on n-alkane by Cladosporium resinae* Appl. Microbiol. Biotechnol., **51**, 860-864
4. Nishida, K., Yagisawa, F., Kuroiwa, H., Nagata, T. and Kuroiwa, T. (2005) *Cell cycle-regulated microtubeli-independent organelle division in Cyanidioschyzon merolae* Mol. Biol. Cell, **16**, 2493-2502
5. Nishida, K., Yagisawa, F., Kuroiwa, H., Yoshida, Y. and Kuroiwa, T. (2007) *WD40 protein Mdal is purified with Dnm1 and forms a dividing ring for mitochondria before Dnm1 in Cyanidioschyzon merolae* Proc. Natl. Acad. Sci. USA, **104**, 4736-4741
6. Yagisawa, F., Nishida, K., Yoshida, M., Ohnuma, M., Shimada, T., Fujiwara, T., Yoshida, Y., Misumi, O., Kuroiwa, H. and Kuroiwa, T. (2009) *Identification of novel proteins in isolated polyphosphate vacuoles in the primitive red alga Cyanidioschyzon merolae* Plant J., **60**, 882-893
7. Imoto, Y., Abe, Y., Okumoto, K., Honsho, M., Kuroiwa, H., Kuroiwa, T. and Fujiki, Y. (2017) *Defining the dynamin-based ring organizing center on the peroxisome-dividing machinery isolated from Cyanidioschyzon merolae* J. Cell Sci., **130**, 853-867
8. Oliveira, D.L., Nimrichter, L., Miranda, K., Frases, S., Faull, K.F., Casadevall, A. and Rodrigues, M.L. (2009) *Cryptococcus neoformans cryoultramicrotomy and vesicle fractionation reveals an intimate association between membrane lipids and glucuronoxylomannan* Fungal Genet. Biol., **46**, 956-963
9. Wolf, J.M., Rivera, J. and Casadevall, A. (2012) *Serum albumin disrupts Cryptococcus neoformans and Bacillus anthracis extracellular vesicles* Cellular Microbiology (2012) 14(5), 762-773
10. Managadze, D., Würtz, C., Wiese, S., Meyer, H.E., Niehaus, G., Erdmann, R., Warscheid, B. and Rottenecker, H. (2010) *A proteomic approach towards the identification of the matrix protein content of the two types of microbodies in Neurospora crassa* Proteomics, **10**, 3222-3234
11. Brito, W.deA., Rezende, T.C.V., Parente, A.F., Ricart, C.A.O., de Sousa, M.V., Bão, N. and Soares, C.M.deA. (2011) *Identification, characterization and regulation studies of the aconitase of Paracoccidioides brasiliensis* Fungal Biol., **115**, 697-707

12. Diener, D.R., Lupetti, P. and Rosenbaum, J.L. (2015) *Proteomic analysis of isolated ciliary transition zones reveals the presence of ESCRT proteins* Curr. Biol., **25**, 379–384
13. Caballero-Lima, D., Hautbergue, G.M., Wilson, S.A. and Sudbery, P.E. (2014) *In Candida albicans hyphae, Sec2p is physically associated with SEC2 mRNA on secretory vesicles* Mol. Microbiol., **94**, 828–842

OptiPrep™ Application Sheet S59; 6th edition, January 2020